

Semi-closed form prices of barrier options in the time-dependent CEV and CIR models

Andrey Itkin

joined work with Peter Carr and Dmitry Muravey

Tandon School of Engineering, NYU

QuantMinds Americas, 2020

Disclaimer

The views represented herein are the author's own views and do not necessarily represent the views of BoA and NYU or their affiliates and are not a product of BoA Research.

The time-dependent constant elasticity of variance (CEV) model is a one-dimensional diffusion process that solves a stochastic differential equation

$$dS_t = \mu(t)S_t dt + \sigma(t)S_t^{\beta+1} dW_t, \quad S_{t=0} = S_0. \quad (1)$$

Here $t \geq 0$ is the time, S_t is the stochastic stock price, $\mu(t)$ is the drift, $\sigma(t)$ is the volatility and β is the elasticity parameter such that $\beta < 1$, $\beta \neq \{0, -1\}$ ¹, W_t is the standard Brownian motion.

¹In case $\beta = 0$ this model is the Black-Scholes model, while for $\beta = -1$ this is the Bachelier, or time-dependent Ornstein-Uhlenbeck (OU) model.

The time-dependent constant elasticity of variance (CEV) model is a one-dimensional diffusion process that solves a stochastic differential equation

$$dS_t = \mu(t)S_t dt + \sigma(t)S_t^{\beta+1} dW_t, \quad S_{t=0} = S_0. \quad (1)$$

Here $t \geq 0$ is the time, S_t is the stochastic stock price, $\mu(t)$ is the drift, $\sigma(t)$ is the volatility and β is the elasticity parameter such that $\beta < 1$, $\beta \neq \{0, -1\}$ ¹, W_t is the standard Brownian motion.

Let us consider an Up-and-Out barrier Call option $C(t, S)$ written on the underlying process S_t . By the Feynman–Kac formula this price solves the following partial differential equation (PDE)

$$\frac{\partial C}{\partial t} + \frac{1}{2}\sigma^2(t)S^{2\beta+2}\frac{\partial^2 C}{\partial S^2} + [r(t) - q(t)]S\frac{\partial C}{\partial S} = r(t)C. \quad (2)$$

¹In case $\beta = 0$ this model is the Black-Scholes model, while for $\beta = -1$ this is the Bachelier, or time-dependent Ornstein-Uhlenbeck (OU) model.

The time-dependent constant elasticity of variance (CEV) model is a one-dimensional diffusion process that solves a stochastic differential equation

$$dS_t = \mu(t)S_t dt + \sigma(t)S_t^{\beta+1} dW_t, \quad S_{t=0} = S_0. \quad (1)$$

Here $t \geq 0$ is the time, S_t is the stochastic stock price, $\mu(t)$ is the drift, $\sigma(t)$ is the volatility and β is the elasticity parameter such that $\beta < 1$, $\beta \neq \{0, -1\}$ ¹, W_t is the standard Brownian motion.

Let us consider an Up-and-Out barrier Call option $C(t, S)$ written on the underlying process S_t . By the Feynman–Kac formula this price solves the following partial differential equation (PDE)

$$\frac{\partial C}{\partial t} + \frac{1}{2}\sigma^2(t)S^{2\beta+2}\frac{\partial^2 C}{\partial S^2} + [r(t) - q(t)]S\frac{\partial C}{\partial S} = r(t)C. \quad (2)$$

This equation should be solved subject to the terminal condition at the option maturity $t = T$

$$C(T, S) = (S - K)^+, \quad (3)$$

where K is the option strike, and the boundary conditions

$$C(t, 0) = 0, \quad C(t, H(t)) = 0, \quad (4)$$

where $H(t)$ is the upper barrier, perhaps time-dependent.

¹In case $\beta = 0$ this model is the Black-Scholes model, while for $\beta = -1$ this is the Bachelier, or time-dependent Ornstein-Uhlenbeck (OU) model.

Proposition

The PDE in Eq.(2) can be transformed to

$$\frac{\partial u}{\partial \tau} = \frac{1}{2} \frac{\partial^2 u}{\partial z^2} + \frac{b}{z} \frac{\partial u}{\partial z}, \quad (5)$$

where b is some constant, $u = u(\tau, z)$ is the new dependent variable, and (τ, z) are the new independent variables. The Eq.(5) is the PDE associated with the one-dimensional Bessel process, [Revuz and Yor 1999]

$$dX_t = dW_t + \frac{b}{X_t} dt. \quad (6)$$

Proposition

The PDE in Eq.(2) can be transformed to

$$\frac{\partial u}{\partial \tau} = \frac{1}{2} \frac{\partial^2 u}{\partial z^2} + \frac{b}{z} \frac{\partial u}{\partial z}, \quad (5)$$

where b is some constant, $u = u(\tau, z)$ is the new dependent variable, and (τ, z) are the new independent variables. The Eq.(5) is the PDE associated with the one-dimensional Bessel process, [Revuz and Yor 1999]

$$dX_t = dW_t + \frac{b}{X_t} dt. \quad (6)$$

One note:

- Carr and Linetsky in [Carr and Linetsky 2006] extend the time-dependent CEV model considered in this paper by allowing a jump to zero. They also reduce their stock price process to a time homogeneous Bessel process.

Proposition

The PDE in Eq.(2) can be transformed to

$$\frac{\partial u}{\partial \tau} = \frac{1}{2} \frac{\partial^2 u}{\partial z^2} + \frac{b}{z} \frac{\partial u}{\partial z}, \quad (5)$$

where b is some constant, $u = u(\tau, z)$ is the new dependent variable, and (τ, z) are the new independent variables. The Eq.(5) is the PDE associated with the one-dimensional Bessel process, [Revuz and Yor 1999]

$$dX_t = dW_t + \frac{b}{X_t} dt. \quad (6)$$

One note:

- Carr and Linetsky in [Carr and Linetsky 2006] extend the time-dependent CEV model considered in this paper by allowing a jump to zero. They also reduce their stock price process to a time homogeneous Bessel process.

As far as the terminal (now the initial) condition in Eq.(3) and the boundary conditions in Eq.(4) in the new variables is concerned, we must distinguish two cases, which are determined by the sign of β .

- If $-1 < \beta < 0$, the domain of definition for z is $z \in [0, y(\tau)]$, where $y(\tau)$ can be explicitly represented in terms of the parameters $\mu(t)$, $\sigma(t)$, β and the barrier $H(t)$.

As far as the terminal (now the initial) condition in Eq.(3) and the boundary conditions in Eq.(4) in the new variables is concerned, we must distinguish two cases, which are determined by the sign of β .

- If $-1 < \beta < 0$, the domain of definition for z is $z \in [0, y(\tau)]$, where $y(\tau)$ can be explicitly represented in terms of the parameters $\mu(t)$, $\sigma(t)$, β and the barrier $H(t)$.
- Therefore, the price of the Up-and-Out barrier Call option solves the following initial-boundary problem in the curvilinear strip $[0, y(\tau))$:

$$\begin{aligned} \frac{\partial u(\tau, z)}{\partial \tau} &= \frac{1}{2} \frac{\partial^2 u(\tau, z)}{\partial z^2} + \frac{b}{u} \frac{\partial u(\tau, z)}{\partial z}, \\ u(0, z) &= f(z), \quad 0 < z < y(0), \\ u(\tau, 0) &= 0, \quad u(\tau, y(\tau)) = 0. \end{aligned} \tag{7}$$

As far as the terminal (now the initial) condition in Eq.(3) and the boundary conditions in Eq.(4) in the new variables is concerned, we must distinguish two cases, which are determined by the sign of β .

- If $-1 < \beta < 0$, the domain of definition for z is $z \in [0, y(\tau)]$, where $y(\tau)$ can be explicitly represented in terms of the parameters $\mu(t)$, $\sigma(t)$, β and the barrier $H(t)$.
- Therefore, the price of the Up-and-Out barrier Call option solves the following initial-boundary problem in the curvilinear strip $[0, y(\tau))$:

$$\begin{aligned}\frac{\partial u(\tau, z)}{\partial \tau} &= \frac{1}{2} \frac{\partial^2 u(\tau, z)}{\partial z^2} + \frac{b}{u} \frac{\partial u(\tau, z)}{\partial z}, \\ u(0, z) &= f(z), \quad 0 < z < y(0), \\ u(\tau, 0) &= 0, \quad u(\tau, y(\tau)) = 0.\end{aligned}\tag{7}$$

- However, if $0 < \beta < 1$, the left boundary goes to $-\infty$. In this case we obtain the problem in the half-plane $[y(\tau), \infty)$:

$$\begin{aligned}\frac{\partial u(\tau, z)}{\partial \tau} &= \frac{1}{2} \frac{\partial^2 u(\tau, z)}{\partial z^2} + \frac{b}{u} \frac{\partial u(\tau, z)}{\partial z}, \\ u(0, z) &= f(z), \quad y(0) < z < \infty, \\ u(\tau, z) \Big|_{z \rightarrow \infty} &= 0, \quad u(\tau, y(\tau)) = 0.\end{aligned}\tag{8}$$

As far as the terminal (now the initial) condition in Eq.(3) and the boundary conditions in Eq.(4) in the new variables is concerned, we must distinguish two cases, which are determined by the sign of β .

- If $-1 < \beta < 0$, the domain of definition for z is $z \in [0, y(\tau)]$, where $y(\tau)$ can be explicitly represented in terms of the parameters $\mu(t)$, $\sigma(t)$, β and the barrier $H(t)$.
- Therefore, the price of the Up-and-Out barrier Call option solves the following initial-boundary problem in the curvilinear strip $[0, y(\tau))$:

$$\begin{aligned}\frac{\partial u(\tau, z)}{\partial \tau} &= \frac{1}{2} \frac{\partial^2 u(\tau, z)}{\partial z^2} + \frac{b}{u} \frac{\partial u(\tau, z)}{\partial z}, \\ u(0, z) &= f(z), \quad 0 < z < y(0), \\ u(\tau, 0) &= 0, \quad u(\tau, y(\tau)) = 0.\end{aligned}\tag{7}$$

- However, if $0 < \beta < 1$, the left boundary goes to $-\infty$. In this case we obtain the problem in the half-plane $[y(\tau), \infty)$:

$$\begin{aligned}\frac{\partial u(\tau, z)}{\partial \tau} &= \frac{1}{2} \frac{\partial^2 u(\tau, z)}{\partial z^2} + \frac{b}{u} \frac{\partial u(\tau, z)}{\partial z}, \\ u(0, z) &= f(z), \quad y(0) < z < \infty, \\ u(\tau, z) \Big|_{z \rightarrow \infty} &= 0, \quad u(\tau, y(\tau)) = 0.\end{aligned}\tag{8}$$

The time-dependent version of the CIR model can be defined as a solution of the following SDE:

$$dr_t = \kappa(t)[\theta(t) - r_t]dt + \sigma(t)\sqrt{r_t}dW_t, \quad r_{t=0} = r. \quad (9)$$

Here $\kappa(t) > 0$ is the mean-reversion speed, and $\theta(t)$ is the mean-reversion level.

The time-dependent version of the CIR model can be defined as a solution of the following SDE:

$$dr_t = \kappa(t)[\theta(t) - r_t]dt + \sigma(t)\sqrt{r_t}dW_t, \quad r_{t=0} = r. \quad (9)$$

Here $\kappa(t) > 0$ is the mean-reversion speed, and $\theta(t)$ is the mean-reversion level.

Let us consider a Down-and-Out barrier Call option written on a ZCB as an underlying. Under a risk-neutral measure the option price $C(t, r)$ solves the following PDE, [Andersen and Piterbarg 2010] .

$$\frac{\partial C}{\partial t} + \frac{1}{2}\sigma^2(t)r\frac{\partial^2 C}{\partial r^2} + \kappa(t)[\theta(t) - r]\frac{\partial C}{\partial r} = rC. \quad (10)$$

The terminal condition at the option maturity $T \leq S$ for this PDE reads

$$C(T, r) = (F(r, T, S) - K)^+, \quad (11)$$

where K is the option strike, $F(r, t, S)$ is the ZCB price, and S is the bond expiration.

The time-dependent version of the CIR model can be defined as a solution of the following SDE:

$$dr_t = \kappa(t)[\theta(t) - r_t]dt + \sigma(t)\sqrt{r_t}dW_t, \quad r_{t=0} = r. \quad (9)$$

Here $\kappa(t) > 0$ is the mean-reversion speed, and $\theta(t)$ is the mean-reversion level.

Let us consider a Down-and-Out barrier Call option written on a ZCB as an underlying. Under a risk-neutral measure the option price $C(t, r)$ solves the following PDE, [Andersen and Piterbarg 2010] .

$$\frac{\partial C}{\partial t} + \frac{1}{2}\sigma^2(t)r\frac{\partial^2 C}{\partial r^2} + \kappa(t)[\theta(t) - r]\frac{\partial C}{\partial r} = rC. \quad (10)$$

The terminal condition at the option maturity $T \leq S$ for this PDE reads

$$C(T, r) = (F(r, T, S) - K)^+, \quad (11)$$

where K is the option strike, $F(r, t, S)$ is the ZCB price, and S is the bond expiration.

The boundary condition is

$$C(t, r) = 0 \quad \text{if } F(r, t, S) = L_F(t). \quad (12)$$

The time-dependent version of the CIR model can be defined as a solution of the following SDE:

$$dr_t = \kappa(t)[\theta(t) - r_t]dt + \sigma(t)\sqrt{r_t}dW_t, \quad r_{t=0} = r. \quad (9)$$

Here $\kappa(t) > 0$ is the mean-reversion speed, and $\theta(t)$ is the mean-reversion level.

Let us consider a Down-and-Out barrier Call option written on a ZCB as an underlying. Under a risk-neutral measure the option price $C(t, r)$ solves the following PDE, [Andersen and Piterbarg 2010] .

$$\frac{\partial C}{\partial t} + \frac{1}{2}\sigma^2(t)r\frac{\partial^2 C}{\partial r^2} + \kappa(t)[\theta(t) - r]\frac{\partial C}{\partial r} = rC. \quad (10)$$

The terminal condition at the option maturity $T \leq S$ for this PDE reads

$$C(T, r) = (F(r, T, S) - K)^+, \quad (11)$$

where K is the option strike, $F(r, t, S)$ is the ZCB price, and S is the bond expiration.

The boundary condition is

$$C(t, r) = 0 \quad \text{if } F(r, t, S) = L_F(t). \quad (12)$$

Since the CIR model belongs to the class of exponentially affine models, the price of the ZCB $F(r, t, S)$ for this model is known in closed form. Thus, this condition can be translated into the r domain ($r = L(t)$).

The time-dependent version of the CIR model can be defined as a solution of the following SDE:

$$dr_t = \kappa(t)[\theta(t) - r_t]dt + \sigma(t)\sqrt{r_t}dW_t, \quad r_{t=0} = r. \quad (9)$$

Here $\kappa(t) > 0$ is the mean-reversion speed, and $\theta(t)$ is the mean-reversion level.

Let us consider a Down-and-Out barrier Call option written on a ZCB as an underlying. Under a risk-neutral measure the option price $C(t, r)$ solves the following PDE, [Andersen and Piterbarg 2010] .

$$\frac{\partial C}{\partial t} + \frac{1}{2}\sigma^2(t)r\frac{\partial^2 C}{\partial r^2} + \kappa(t)[\theta(t) - r]\frac{\partial C}{\partial r} = rC. \quad (10)$$

The terminal condition at the option maturity $T \leq S$ for this PDE reads

$$C(T, r) = (F(r, T, S) - K)^+, \quad (11)$$

where K is the option strike, $F(r, t, S)$ is the ZCB price, and S is the bond expiration.

The boundary condition is

$$C(t, r) = 0 \quad \text{if } F(r, t, S) = L_F(t). \quad (12)$$

Since the CIR model belongs to the class of exponentially affine models, the price of the ZCB $F(r, t, S)$ for this model is known in closed form. Thus, this condition can be translated into the r domain ($r = L(t)$).

The second boundary can be naturally set at $r \rightarrow \infty$. As at $r \rightarrow \infty$ the ZCB price tends to zero, the Call option price also vanishes in this limit.

Proposition

The Eq.(10) can be transformed to

$$\frac{\partial u}{\partial \tau} = \frac{1}{2} \frac{\partial^2 u}{\partial z^2} + \frac{b}{z} \frac{\partial u}{\partial z}, \quad (13)$$

where b is some constant, $u = u(\tau, z)$ is the new dependent variable, and (τ, z) are the new independent variables, if

$$\frac{\kappa(t)\theta(t)}{\sigma^2(t)} = \frac{m}{2}, \quad (14)$$

where $m \in [0, \infty)$ is some constant.

Proposition

The Eq.(10) can be transformed to

$$\frac{\partial u}{\partial \tau} = \frac{1}{2} \frac{\partial^2 u}{\partial z^2} + \frac{b}{z} \frac{\partial u}{\partial z}, \quad (13)$$

where b is some constant, $u = u(\tau, z)$ is the new dependent variable, and (τ, z) are the new independent variables, if

$$\frac{\kappa(t)\theta(t)}{\sigma^2(t)} = \frac{m}{2}, \quad (14)$$

where $m \in [0, \infty)$ is some constant.

Two notes:

- The transformation for the time-dependent CIR model from Eq.(10) to Eq.(13) cannot be done unconditionally.

Proposition

The Eq.(10) can be transformed to

$$\frac{\partial u}{\partial \tau} = \frac{1}{2} \frac{\partial^2 u}{\partial z^2} + \frac{b}{z} \frac{\partial u}{\partial z}, \quad (13)$$

where b is some constant, $u = u(\tau, z)$ is the new dependent variable, and (τ, z) are the new independent variables, if

$$\frac{\kappa(t)\theta(t)}{\sigma^2(t)} = \frac{m}{2}, \quad (14)$$

where $m \in [0, \infty)$ is some constant.

Two notes:

- ▶ The transformation for the time-dependent CIR model from Eq.(10) to Eq.(13) cannot be done unconditionally.
- ▶ However, from practitioners' points of view the condition Eq.(14) seems not to be too restrictive. Since m is an arbitrary constant, it could be calibrated to the market data together with $\kappa(t)$ and $\sigma(t)$. Therefore, in this form the model should be capable for calibration to the term-structure of interest rates.

Proposition

The Eq.(10) can be transformed to

$$\frac{\partial u}{\partial \tau} = \frac{1}{2} \frac{\partial^2 u}{\partial z^2} + \frac{b}{z} \frac{\partial u}{\partial z}, \quad (13)$$

where b is some constant, $u = u(\tau, z)$ is the new dependent variable, and (τ, z) are the new independent variables, if

$$\frac{\kappa(t)\theta(t)}{\sigma^2(t)} = \frac{m}{2}, \quad (14)$$

where $m \in [0, \infty)$ is some constant.

Two notes:

- ▶ The transformation for the time-dependent CIR model from Eq.(10) to Eq.(13) cannot be done unconditionally.
- ▶ However, from practitioners' points of view the condition Eq.(14) seems not to be too restrictive. Since m is an arbitrary constant, it could be calibrated to the market data together with $\kappa(t)$ and $\sigma(t)$. Therefore, in this form the model should be capable for calibration to the term-structure of interest rates.

We start the exposition of the method of Bessel potentials for the problem in the half-plane $[y(\tau), \infty)$.

We start the exposition of the method of Bessel potentials for the problem in the half-plane $[y(\tau), \infty)$.

Since both the CEV and CIR problems have the inhomogeneous initial condition, our first step is to reduce them to the alternative problems with a homogeneous initial condition. The Green function of the Bessel equation at the infinite domain is known in closed form, [Polyanin 2002], therefore this can be achieved by representing $u(\tau, z)$ in the form

$$u(\tau, z) = q(\tau, z) + \int_{y(0)}^{\infty} u(0, \zeta) q_{\tau}(z, \zeta, b) d\zeta. \quad (15)$$

We start the exposition of the method of Bessel potentials for the problem in the half-plane $[y(\tau), \infty)$.

Since both the CEV and CIR problems have the inhomogeneous initial condition, our first step is to reduce them to the alternative problems with a homogeneous initial condition. The Green function of the Bessel equation at the infinite domain is known in closed form, [Polyanin 2002], therefore this can be achieved by representing $u(\tau, z)$ in the form

$$u(\tau, z) = q(\tau, z) + \int_{y(0)}^{\infty} u(0, \zeta) q_{\tau}(z, \zeta, b) d\zeta. \quad (15)$$

Here $q_{\tau}(z, \zeta, b)$ is the fundamental solution (or the transition density, or the Green function) of Eq.(13) at the domain $z \in [0, \infty)$.

We start the exposition of the method of Bessel potentials for the problem in the half-plane $[y(\tau), \infty)$.

Since both the CEV and CIR problems have the inhomogeneous initial condition, our first step is to reduce them to the alternative problems with a homogeneous initial condition. The Green function of the Bessel equation at the infinite domain is known in closed form, [Polyanin 2002], therefore this can be achieved by representing $u(\tau, z)$ in the form

$$u(\tau, z) = q(\tau, z) + \int_{y(0)}^{\infty} u(0, \zeta) q_{\tau}(z, \zeta, b) d\zeta. \quad (15)$$

Here $q_{\tau}(z, \zeta, b)$ is the fundamental solution (or the transition density, or the Green function) of Eq.(13) at the domain $z \in [0, \infty)$.

For the CEV model $b = (1 + \beta) / (2\beta)$. Since in this case $0 < \beta < 1$, we get $b > 1$. For the CIR model $b = m - 1/2$ where $m \in [0, \infty)$ can be found by calibration. It is known, [Lawler 2018; Linetsky and Mendoza 2010], that in case $b \geq 1/2$ the density $q_{\tau}(z, \zeta, b)$ is a good density with no defect of mass, i.e., it integrates into 1.

We start the exposition of the method of Bessel potentials for the problem in the half-plane $[y(\tau), \infty)$.

Since both the CEV and CIR problems have the inhomogeneous initial condition, our first step is to reduce them to the alternative problems with a homogeneous initial condition. The Green function of the Bessel equation at the infinite domain is known in closed form, [Polyanin 2002], therefore this can be achieved by representing $u(\tau, z)$ in the form

$$u(\tau, z) = q(\tau, z) + \int_{y(0)}^{\infty} u(0, \zeta) q_{\tau}(z, \zeta, b) d\zeta. \quad (15)$$

Here $q_{\tau}(z, \zeta, b)$ is the fundamental solution (or the transition density, or the Green function) of Eq.(13) at the domain $z \in [0, \infty)$.

For the CEV model $b = (1 + \beta) / (2\beta)$. Since in this case $0 < \beta < 1$, we get $b > 1$. For the CIR model $b = m - 1/2$ where $m \in [0, \infty)$ can be found by calibration. It is known, [Lawler 2018; Linetsky and Mendoza 2010], that in case $b \geq 1/2$ the density $q_{\tau}(z, \zeta, b)$ is a good density with no defect of mass, i.e., it integrates into 1. The explicit representation reads, [Cox 1975; Emanuel and Macbeth 1982]

$$q_{\tau}(z, \zeta, b) = \frac{\sqrt{z\zeta}}{\tau} \left(\frac{\zeta}{z}\right)^b e^{-\frac{z^2 + \zeta^2}{2\tau}} I_{b-1/2} \left(\frac{z\zeta}{\tau}\right). \quad (16)$$

Here $I_{\nu}(x)$ is the modified Bessel function of the first kind, [Abramowitz and Stegun 1964].

The function $q(x, \tau)$ solves the problem with zero initial condition:

$$\begin{aligned}
 \frac{\partial q(\tau, z)}{\partial \tau} &= \frac{1}{2} \frac{\partial^2 q(\tau, z)}{\partial z^2} + \frac{b}{z} \frac{\partial q(\tau, z)}{\partial z}, \\
 q(0, z) &= 0, \quad y(0) < z < \infty, \\
 q(\tau, z) \Big|_{z \rightarrow \infty} &= 0, \quad q(\tau, y(\tau)) = \zeta(\tau), \\
 \zeta(\tau) &= - \int_{y(0)}^{\infty} u(0, \zeta) q_{\tau}(y(\tau), \zeta, b) d\zeta.
 \end{aligned} \tag{17}$$

The function $q(x, \tau)$ solves the problem with zero initial condition:

$$\begin{aligned} \frac{\partial q(\tau, z)}{\partial \tau} &= \frac{1}{2} \frac{\partial^2 q(\tau, z)}{\partial z^2} + \frac{b}{z} \frac{\partial q(\tau, z)}{\partial z}, \\ q(0, z) &= 0, \quad y(0) < z < \infty, \\ q(\tau, z) \Big|_{z \rightarrow \infty} &= 0, \quad q(\tau, y(\tau)) = \zeta(\tau), \\ \zeta(\tau) &= - \int_{y(0)}^{\infty} u(0, \zeta) q_{\tau}(y(\tau), \zeta, b) d\zeta. \end{aligned} \quad (17)$$

Therefore, following the general idea of the method of heat potentials, we represent the solution in the form of a generalized potential for the Bessel PDE

$$q(\tau, z) = \int_0^{\tau} \Psi(k) \frac{\partial}{\partial \tilde{\zeta}} \left[\frac{\sqrt{z\tilde{\zeta}}}{\tau - k} \left(\frac{\tilde{\zeta}}{z} \right)^b e^{-\frac{z^2 + \tilde{\zeta}^2}{2(\tau - k)}} I_{b-1/2} \left(\frac{z\tilde{\zeta}}{\tau - k} \right) \right] \Big|_{\tilde{\zeta} \rightarrow y(k)} dk, \quad (18)$$

where $\Psi(k)$ is the potential density.

The function $q(x, \tau)$ solves the problem with zero initial condition:

$$\begin{aligned} \frac{\partial q(\tau, z)}{\partial \tau} &= \frac{1}{2} \frac{\partial^2 q(\tau, z)}{\partial z^2} + \frac{b}{z} \frac{\partial q(\tau, z)}{\partial z}, \\ q(0, z) &= 0, \quad y(0) < z < \infty, \\ q(\tau, z) \Big|_{z \rightarrow \infty} &= 0, \quad q(\tau, y(\tau)) = \zeta(\tau), \\ \zeta(\tau) &= - \int_{y(0)}^{\infty} u(0, \zeta) q_{\tau}(y(\tau), \zeta, b) d\zeta. \end{aligned} \quad (17)$$

Therefore, following the general idea of the method of heat potentials, we represent the solution in the form of a generalized potential for the Bessel PDE

$$q(\tau, z) = \int_0^{\tau} \Psi(k) \frac{\partial}{\partial \tilde{\zeta}} \left[\frac{\sqrt{z\tilde{\zeta}}}{\tau - k} \left(\frac{\tilde{\zeta}}{z} \right)^b e^{-\frac{z^2 + \tilde{\zeta}^2}{2(\tau - k)}} I_{b-1/2} \left(\frac{z\tilde{\zeta}}{\tau - k} \right) \right] \Big|_{\tilde{\zeta} \rightarrow y(k)} dk, \quad (18)$$

where $\Psi(k)$ is the potential density.

It can be seen that $q(\tau, z)$ solves Eq.(13) as the derivative of the integral on the upper limit is proportional to the Delta function which vanishes due to $z \neq y(\tau)$. The solution in Eq.(18) also satisfies the initial condition at $\tau = 0$, and the vanishing condition at $z \rightarrow \infty$.

At the barrier $z = y(\tau)$ function $q(\tau, z)$ is discontinuous. Following a similar approach for the heat potentials method, [Tikhonov and Samarskii 1963]), it can be shown that the limiting value of $q(\tau, z)$ at $z = y(\tau) + 0$ is equal to $\zeta(\tau)$:

$$\zeta(\tau) = \Psi(\tau) + \int_0^\tau \Psi(k) \frac{\partial}{\partial y(k)} \left[\frac{\sqrt{y(\tau)y(k)}}{\tau - k} \left(\frac{y(k)}{y(\tau)} \right)^b e^{-\frac{y^2(\tau) + y^2(k)}{2(\tau - k)}} I_{b-1/2} \left(\frac{y(\tau)y(k)}{\tau - k} \right) \right] dk. \quad (19)$$

At the barrier $z = y(\tau)$ function $q(\tau, z)$ is discontinuous. Following a similar approach for the heat potentials method, [Tikhonov and Samarskii 1963]), it can be shown that the limiting value of $q(\tau, z)$ at $z = y(\tau) + 0$ is equal to $\zeta(\tau)$:

$$\zeta(\tau) = \Psi(\tau) + \int_0^\tau \Psi(k) \frac{\partial}{\partial y(k)} \left[\frac{\sqrt{y(\tau)y(k)}}{\tau - k} \left(\frac{y(k)}{y(\tau)} \right)^b e^{-\frac{y^2(\tau) + y^2(k)}{2(\tau - k)}} I_{b-1/2} \left(\frac{y(\tau)y(k)}{\tau - k} \right) \right] dk. \quad (19)$$

The Eq.(19) is a linear Volterra equations of the second kind, [Polyanin and Manzhirov 2008] . Since $\zeta(\tau)$ is a continuously differentiable function, Eq.(19) has a unique continuous solution for $\Psi(\tau)$. The Volterra equation can be efficiently solved either numerically or semi-analytically, see [Itkin and Muravey 2020] for the discussion on various approaches to solving this type of equations.

At the barrier $z = y(\tau)$ function $q(\tau, z)$ is discontinuous. Following a similar approach for the heat potentials method, [Tikhonov and Samarskii 1963]), it can be shown that the limiting value of $q(\tau, z)$ at $z = y(\tau) + 0$ is equal to $\zeta(\tau)$:

$$\zeta(\tau) = \Psi(\tau) + \int_0^\tau \Psi(k) \frac{\partial}{\partial y(k)} \left[\frac{\sqrt{y(\tau)y(k)}}{\tau - k} \left(\frac{y(k)}{y(\tau)} \right)^b e^{-\frac{y^2(\tau) + y^2(k)}{2(\tau - k)}} I_{b-1/2} \left(\frac{y(\tau)y(k)}{\tau - k} \right) \right] dk. \quad (19)$$

The Eq.(19) is a linear Volterra equations of the second kind, [Polyanin and Manzhirov 2008] . Since $\zeta(\tau)$ is a continuously differentiable function, Eq.(19) has a unique continuous solution for $\Psi(\tau)$. The Volterra equation can be efficiently solved either numerically or semi-analytically, see [Itkin and Muravey 2020] for the discussion on various approaches to solving this type of equations.

Once Eq.(19) is solved and the function $\Psi(\tau)$ is found, the final solution reads

$$u(\tau, z) = \int_0^\tau \Psi(k) \frac{\partial}{\partial y(k)} \left[\frac{\sqrt{zy(k)}}{\tau - k} \left(\frac{y(k)}{z} \right)^b e^{-\frac{z^2 + y^2(k)}{2(\tau - k)}} I_{b-1/2} \left(\frac{zy(k)}{\tau - k} \right) \right] dk + \int_{y(0)}^\infty u(0, \zeta) q_\tau(z, \zeta, b) d\zeta. \quad (20)$$

Consider now the problem in the domain $[0, y(\tau)]$. The construction of the solution is similar to the presented above. Again, in order to obtain a PDE with a homogeneous initial condition, we represent the solution in the form

$$u(\tau, z) = q(\tau, z) - \varsigma_0(\tau) + \int_0^{y(0)} u(0, \xi) q_\tau(z, \xi, b) d\xi, \quad (21)$$

$$\varsigma_0(\tau) = - \int_0^{y(0)} u(0, \xi) q_\tau(0, \xi, b) d\xi,$$

$$q_\tau(0, \xi, b) = \frac{2^{1/2-b} \xi^{2b}}{\tau^{b+1/2} \Gamma(b + \frac{1}{2})} e^{-\frac{\xi^2}{2\tau}},$$

where ($\Gamma(x)$ is the Euler Gamma function, [Abramowitz and Stegun 1964]). The method to construct $q(\tau, z)$ is fully analogical to the previous section.

Consider now the problem in the domain $[0, y(\tau)]$. The construction of the solution is similar to the presented above. Again, in order to obtain a PDE with a homogeneous initial condition, we represent the solution in the form

$$u(\tau, z) = q(\tau, z) - \varsigma_0(\tau) + \int_0^{y(0)} u(0, \xi) q_\tau(z, \xi, b) d\xi, \quad (21)$$

$$\varsigma_0(\tau) = - \int_0^{y(0)} u(0, \xi) q_\tau(0, \xi, b) d\xi,$$

$$q_\tau(0, \xi, b) = \frac{2^{1/2-b} \xi^{2b}}{\tau^{b+1/2} \Gamma(b + \frac{1}{2})} e^{-\frac{\xi^2}{2\tau}},$$

where ($\Gamma(x)$ is the Euler Gamma function, [Abramowitz and Stegun 1964]). The method to construct $q(\tau, z)$ is fully analogical to the previous section.

The double barrier options can also be priced via the BP method. This problem corresponds to the domain $[y(\tau), h(\tau)]$. For doing so, we have to use the so-called double layer potentials:

$$u(\tau, z) = q_y(\tau, z) + q_h(\tau, z) + \int_{y(0)}^{h(0)} u(0, \xi) q_\tau(z, \xi, b) d\xi. \quad (22)$$

However, for this case there are two unknown potential density functions $\Psi(\tau)$ and $\Phi(\tau)$ defined as a solution to a system of Volterra equation. For more details, see [Carr, Itkin, and Muravey 2020].

In this section we describe the method of Generalized integral transform(GIT). We consider the problem in the curvilinear strip $[0, y(\tau)]$. This problem emerges when we consider the CEV process with $\beta < 0$. Since the Laplace transform of Eq.(13) with time-homogeneous coefficients gives rise to the Bessel ODE, [Abramowitz and Stegun 1964] , it would be natural seeking for the general integral transform in the class of Bessel functions.

In this section we describe the method of Generalized integral transform(GIT). We consider the problem in the curvilinear strip $[0, y(\tau)]$. This problem emerges when we consider the CEV process with $\beta < 0$. Since the Laplace transform of Eq.(13) with time-homogeneous coefficients gives rise to the Bessel ODE, [Abramowitz and Stegun 1964], it would be natural seeking for the general integral transform in the class of Bessel functions. We introduce the GIT of the form

$$\bar{u}(\tau, p) = \int_0^{y(\tau)} z^{\nu+1} u(\tau, z) J_{|\nu|}(zp) dz, \quad (23)$$

where $p = a + i\omega$ is a complex number, $J_\nu(x)$ is the Bessel function of the first kind, and $\nu = 1/(2\beta) < 0$, since $\beta < 0$.

In this section we describe the method of Generalized integral transform (GIT). We consider the problem in the curvilinear strip $[0, y(\tau)]$. This problem emerges when we consider the CEV process with $\beta < 0$. Since the Laplace transform of Eq.(13) with time-homogeneous coefficients gives rise to the Bessel ODE, [Abramowitz and Stegun 1964], it would be natural seeking for the general integral transform in the class of Bessel functions. We introduce the GIT of the form

$$\bar{u}(\tau, p) = \int_0^{y(\tau)} z^{\nu+1} u(\tau, z) J_{|\nu|}(zp) dz, \quad (23)$$

where $p = a + i\omega$ is a complex number, $J_\nu(x)$ is the Bessel function of the first kind, and $\nu = 1/(2\beta) < 0$, since $\beta < 0$.

Next, let us multiply both parts of Eq.(13) by $z^{\nu+1} J_{|\nu|}(zp)$ and integrate on z from 0 to $y(\tau)$. In the result we obtain the following Cauchy problem for \bar{u} (the unknown function $\Psi(\tau)$ should also be found):

$$\begin{aligned} \frac{d\bar{u}(\tau, p)}{d\tau} &= \frac{1}{2} \left[-p^2 \bar{u}(\tau, p) + y^{\nu+1}(\tau) J_{|\nu|}(y(\tau)p) \Psi(\tau) \right], \\ \bar{u}(p, 0) &= \int_0^{y(0)} z^{\nu+1} J_{|\nu|}(zp) u(0, z) dz, \quad \Psi(\tau) = \left. \frac{\partial u}{\partial z} \right|_{z=y(\tau)}. \end{aligned} \quad (24)$$

In this section we describe the method of Generalized integral transform (GIT). We consider the problem in the curvilinear strip $[0, y(\tau)]$. This problem emerges when we consider the CEV process with $\beta < 0$. Since the Laplace transform of Eq.(13) with time-homogeneous coefficients gives rise to the Bessel ODE, [Abramowitz and Stegun 1964], it would be natural seeking for the general integral transform in the class of Bessel functions. We introduce the GIT of the form

$$\bar{u}(\tau, p) = \int_0^{y(\tau)} z^{\nu+1} u(\tau, z) J_{|\nu|}(zp) dz, \quad (23)$$

where $p = a + i\omega$ is a complex number, $J_\nu(x)$ is the Bessel function of the first kind, and $\nu = 1/(2\beta) < 0$, since $\beta < 0$.

Next, let us multiply both parts of Eq.(13) by $z^{\nu+1} J_{|\nu|}(zp)$ and integrate on z from 0 to $y(\tau)$. In the result we obtain the following Cauchy problem for \bar{u} (the unknown function $\Psi(\tau)$ should also be found):

$$\begin{aligned} \frac{d\bar{u}(\tau, p)}{d\tau} &= \frac{1}{2} \left[-p^2 \bar{u}(\tau, p) + y^{\nu+1}(\tau) J_{|\nu|}(y(\tau)p) \Psi(\tau) \right], \\ \bar{u}(p, 0) &= \int_0^{y(0)} z^{\nu+1} J_{|\nu|}(zp) u(0, z) dz, \quad \Psi(\tau) = \left. \frac{\partial u}{\partial z} \right|_{z=y(\tau)}. \end{aligned} \quad (24)$$

The solution of this problem reads

$$\bar{u} = e^{-p^2 \tau / 2} \left[\bar{u}(0, p) + \frac{1}{2} \int_0^\tau e^{p^2 s / 2} \Psi(s) y^{\nu+1}(s) J_{|\nu|}(y(s)p) ds \right]. \quad (25)$$

Note that if $y(\tau) \equiv 0$ the integral transform Eq.(23) is the well known Hankel transform [Bateman and Erdélyi 1953] :

$$u(\tau, z) = z^{-\nu} \int_0^\infty p J_{|\nu|}(zp) \left[\int_0^\infty \zeta^{\nu+1} J_{|\nu|}(\zeta p) u(\tau, \zeta) dz \right] dp \quad (26)$$

:

Note that if $y(\tau) \equiv 0$ the integral transform Eq.(23) is the well known Hankel transform [Bateman and Erdélyi 1953] :

$$u(\tau, z) = z^{-\nu} \int_0^\infty p J_{|\nu|}(zp) \left[\int_0^\infty \zeta^{\nu+1} J_{|\nu|}(\zeta p) u(\tau, \zeta) dz \right] dp \quad (26)$$

Simplifications in the case $y(\tau) = 0$:

- The function Ψ is equal to zero, therefore the Cauchy problem Eq.(24) can be solved explicitly. The inversion formula is known.

:

Note that if $y(\tau) \equiv 0$ the integral transform Eq.(23) is the well known Hankel transform [Bateman and Erdélyi 1953] :

$$u(\tau, z) = z^{-\nu} \int_0^\infty p J_{|\nu|}(zp) \left[\int_0^\infty \zeta^{\nu+1} J_{|\nu|}(\zeta p) u(\tau, \zeta) dz \right] dp \quad (26)$$

Simplifications in the case $y(\tau) = 0$:

- The function Ψ is equal to zero, therefore the Cauchy problem Eq.(24) can be solved explicitly. The inversion formula is known.

:

Note that if $y(\tau) \equiv 0$ the integral transform Eq.(23) is the well known Hankel transform [Bateman and Erdélyi 1953] :

$$u(\tau, z) = z^{-\nu} \int_0^\infty p J_{|\nu|}(zp) \left[\int_0^\infty \zeta^{\nu+1} J_{|\nu|}(\zeta p) u(\tau, \zeta) dz \right] dp \quad (26)$$

Simplifications in the case $y(\tau) = 0$:

- The function Ψ is equal to zero, therefore the Cauchy problem Eq.(24) can be solved explicitly. The inversion formula is known.

The problems arising when using the GIT Eq.(23):

- Problem 1. We have to derive the inversion formula.

Note that if $y(\tau) \equiv 0$ the integral transform Eq.(23) is the well known Hankel transform [Bateman and Erdélyi 1953] :

$$u(\tau, z) = z^{-\nu} \int_0^\infty p J_{|\nu|}(zp) \left[\int_0^\infty \zeta^{\nu+1} J_{|\nu|}(\zeta p) u(\tau, \zeta) dz \right] dp \quad (26)$$

Simplifications in the case $y(\tau) = 0$:

- The function Ψ is equal to zero, therefore the Cauchy problem Eq.(24) can be solved explicitly. The inversion formula is known.

The problems arising when using the GIT Eq.(23):

- Problem 1. We have to derive the inversion formula.
- Problem 2. We need to find the function Ψ .

Note that if $y(\tau) \equiv 0$ the integral transform Eq.(23) is the well known Hankel transform [Bateman and Erdélyi 1953] :

$$u(\tau, z) = z^{-\nu} \int_0^\infty p J_{|\nu|}(zp) \left[\int_0^\infty \zeta^{\nu+1} J_{|\nu|}(\zeta p) u(\tau, \zeta) dz \right] dp \quad (26)$$

Simplifications in the case $y(\tau) = 0$:

- The function Ψ is equal to zero, therefore the Cauchy problem Eq.(24) can be solved explicitly. The inversion formula is known.

The problems arising when using the GIT Eq.(23):

- ▶ Problem 1. We have to derive the inversion formula.
- ▶ Problem 2. We need to find the function Ψ .

Note that if $y(\tau) \equiv 0$ the integral transform Eq.(23) is the well known Hankel transform [Bateman and Erdélyi 1953] :

$$u(\tau, z) = z^{-\nu} \int_0^\infty p J_{|\nu|}(zp) \left[\int_0^\infty \zeta^{\nu+1} J_{|\nu|}(\zeta p) u(\tau, \zeta) dz \right] dp \quad (26)$$

Simplifications in the case $y(\tau) = 0$:

- The function Ψ is equal to zero, therefore the Cauchy problem Eq.(24) can be solved explicitly. The inversion formula is known.

The problems arising when using the GIT Eq.(23):

- ▶ Problem 1. We have to derive the inversion formula.
- ▶ Problem 2. We need to find the function Ψ .

Once these issues are resolved, we obtain a semi-analytical solution to the problem of pricing Up-and-Out barrier Call options for the CEV model with $\beta < 0$.

- ▶ The function $\Psi(\tau)$ can be found as a solution to the Volterra integral equation of the second kind, for more details see [Carr, Itkin, and Muravey 2020] .

Note that if $y(\tau) \equiv 0$ the integral transform Eq.(23) is the well known Hankel transform [Bateman and Erdélyi 1953] :

$$u(\tau, z) = z^{-\nu} \int_0^{\infty} p J_{|\nu|}(zp) \left[\int_0^{\infty} \zeta^{\nu+1} J_{|\nu|}(\zeta p) u(\tau, \zeta) d\zeta \right] dp \quad (26)$$

Simplifications in the case $y(\tau) = 0$:

- The function Ψ is equal to zero, therefore the Cauchy problem Eq.(24) can be solved explicitly. The inversion formula is known.

The problems arising when using the GIT Eq.(23):

- Problem 1. We have to derive the inversion formula.
- Problem 2. We need to find the function Ψ .

Once these issues are resolved, we obtain a semi-analytical solution to the problem of pricing Up-and-Out barrier Call options for the CEV model with $\beta < 0$.

- The function $\Psi(\tau)$ can be found as a solution to the Volterra integral equation of the second kind, for more details see [Carr, Itkin, and Muravey 2020] .
- To build the inverse transform we are looking for the function $u(\tau, z)$ in the form :

$$u(\tau, z) = z^{-\nu} \sum_{n=1}^{\infty} a_n(\tau) J_{|\nu|} \left(\frac{\mu_n z}{y(\tau)} \right). \quad (27)$$

Here μ_n is an ordered sequence of the positive zeros of the Bessel function $J_{|\nu|}(\mu)$:

$$J_{|\nu|}(\mu_n) = J_{|\nu|}(\mu_m) = 0, \quad \mu_n > \mu_m > 0, \quad n > m.$$

Note that if $y(\tau) \equiv 0$ the integral transform Eq.(23) is the well known Hankel transform [Bateman and Erdélyi 1953] :

$$u(\tau, z) = z^{-\nu} \int_0^{\infty} p J_{|\nu|}(zp) \left[\int_0^{\infty} \zeta^{\nu+1} J_{|\nu|}(\zeta p) u(\tau, \zeta) d\zeta \right] dp \quad (26)$$

Simplifications in the case $y(\tau) = 0$:

- The function Ψ is equal to zero, therefore the Cauchy problem Eq.(24) can be solved explicitly. The inversion formula is known.

The problems arising when using the GIT Eq.(23):

- Problem 1. We have to derive the inversion formula.
- Problem 2. We need to find the function Ψ .

Once these issues are resolved, we obtain a semi-analytical solution to the problem of pricing Up-and-Out barrier Call options for the CEV model with $\beta < 0$.

- The function $\Psi(\tau)$ can be found as a solution to the Volterra integral equation of the second kind, for more details see [Carr, Itkin, and Muravey 2020] .
- To build the inverse transform we are looking for the function $u(\tau, z)$ in the form :

$$u(\tau, z) = z^{-\nu} \sum_{n=1}^{\infty} \alpha_n(\tau) J_{|\nu|} \left(\frac{\mu_n z}{y(\tau)} \right). \quad (27)$$

Here μ_n is an ordered sequence of the positive zeros of the Bessel function $J_{|\nu|}(\mu)$:

$$J_{|\nu|}(\mu_n) = J_{|\nu|}(\mu_m) = 0, \quad \mu_n > \mu_m > 0, \quad n > m.$$

Note, that the definition in Eq.(27) automatically respects the vanishing boundary conditions for $u(\tau, z)$. We assume that this series converges absolutely and uniformly $\forall z \in [0, y(\tau)]$ for any $\tau > 0$.

Applying the direct integral transform in Eq.(23) to both parts of Eq.(27), and using a change of variables $z \rightarrow \hat{z} = zy(\tau)$ yields

$$\frac{\bar{u}(\tau, p)}{y^2(\tau)} = \sum_{n=1}^{\infty} \alpha_n(\tau) \int_0^1 \hat{z} J_{|v|}(\mu_n \hat{z}) J_{|v|}(p y(\tau) \hat{z}) d\hat{z}. \quad (28)$$

The set of functions $J_{|v|}(\alpha \hat{z})$ with $\alpha \in \mu_n$, $n = 1, \dots$, forms an orthogonal basis in the space $C[0, 1]$ with the scalar product

$$\langle J_{|v|}(\alpha z), J_{|v|}(\beta z) \rangle = 2 \int_0^1 \frac{z J_{|v|}(\alpha z) J_{|v|}(\beta z) dz}{J_{|v|+1}(\alpha) J_{|v|+1}(\beta)} = \begin{cases} 1, & \alpha = \beta, \\ 0, & \alpha \neq \beta \end{cases} \quad (29)$$

Therefore, the explicit formula for each coefficient $\alpha_n(\tau)$ is straightforward

$$\alpha_n(\tau) = 2 \frac{\bar{u}(\tau, \mu_n / y(\tau))}{y^2(\tau) J_{|v|+1}^2(\mu_n)}. \quad (30)$$

Then the final solution for $u(\tau, z)$ reads

$$\begin{aligned}
 u(\tau, z) = & 2 \frac{z^{-\nu}}{y^2(\tau)} \sum_{n=1}^{\infty} \left[\int_0^{y(0)} u(0, s) s^{\nu+1} e^{-\frac{\mu_n^2}{2y^2(\tau)} \tau} \frac{J_{|\nu|}(\mu_n s / y(\tau)) J_{|\nu|}(\mu_n z / y(\tau))}{J_{|\nu|+1}^2(\mu_n)} ds \right. \\
 & \left. + \frac{1}{2} \int_0^{\tau} y^{\nu+1}(s) \Psi(s) e^{-\frac{\mu_n^2}{2y^2(\tau)} (\tau-s)} \frac{J_{|\nu|}(\mu_n y(s) / y(\tau)) J_{|\nu|}(\mu_n z / y(\tau))}{J_{|\nu|+1}^2(\mu_n)} ds \right].
 \end{aligned} \tag{31}$$

Then the final solution for $u(\tau, z)$ reads

$$\begin{aligned}
 u(\tau, z) = & 2 \frac{z^{-\nu}}{y^2(\tau)} \sum_{n=1}^{\infty} \left[\int_0^{y(0)} u(0, s) s^{\nu+1} e^{-\frac{\mu_n^2}{2y^2(\tau)} \tau} \frac{J_{|\nu|}(\mu_n s / y(\tau)) J_{|\nu|}(\mu_n z / y(\tau))}{J_{|\nu|+1}^2(\mu_n)} ds \right. \\
 & \left. + \frac{1}{2} \int_0^{\tau} y^{\nu+1}(s) \Psi(s) e^{-\frac{\mu_n^2}{2y^2(\tau)} (\tau-s)} \frac{J_{|\nu|}(\mu_n y(s) / y(\tau)) J_{|\nu|}(\mu_n z / y(\tau))}{J_{|\nu|+1}^2(\mu_n)} ds \right]. \quad (31)
 \end{aligned}$$

To obtain the equation for Ψ , one needs to differentiate Eq.(31) on z , and then let $z = y(\tau)$. This yields

$$\begin{aligned}
 \Psi(\tau) = & -\frac{2}{y^{3+\nu}(\tau)} \sum_{n=1}^{\infty} (\mu_n + \nu) \left[\int_0^{y(0)} u(0, s) s^{\nu+1} e^{-\frac{\mu_n^2}{2y^2(\tau)} \tau} \frac{J_{|\nu|}(\mu_n s / y(\tau))}{J_{|\nu|+1}(\mu_n)} ds \right. \\
 & \left. + \frac{1}{2} \int_0^{\tau} y^{\nu+1}(s) \Psi(s) e^{-\frac{\mu_n^2}{2y^2(\tau)} (\tau-s)} \frac{J_{|\nu|}(\mu_n y(s) / y(\tau))}{J_{|\nu|+1}(\mu_n)} ds \right]. \quad (32)
 \end{aligned}$$

Then the final solution for $u(\tau, z)$ reads

$$\begin{aligned}
 u(\tau, z) = & 2 \frac{z^{-\nu}}{y^2(\tau)} \sum_{n=1}^{\infty} \left[\int_0^{y(0)} u(0, s) s^{\nu+1} e^{-\frac{\mu_n^2}{2y^2(\tau)} \tau} \frac{J_{|\nu|}(\mu_n s / y(\tau)) J_{|\nu|}(\mu_n z / y(\tau))}{J_{|\nu|+1}^2(\mu_n)} ds \right. \\
 & \left. + \frac{1}{2} \int_0^{\tau} y^{\nu+1}(s) \Psi(s) e^{-\frac{\mu_n^2}{2y^2(\tau)} (\tau-s)} \frac{J_{|\nu|}(\mu_n y(s) / y(\tau)) J_{|\nu|}(\mu_n z / y(\tau))}{J_{|\nu|+1}^2(\mu_n)} ds \right]. \quad (31)
 \end{aligned}$$

To obtain the equation for Ψ , one needs to differentiate Eq.(31) on z , and then let $z = y(\tau)$. This yields

$$\begin{aligned}
 \Psi(\tau) = & -\frac{2}{y^{3+\nu}(\tau)} \sum_{n=1}^{\infty} (\mu_n + \nu) \left[\int_0^{y(0)} u(0, s) s^{\nu+1} e^{-\frac{\mu_n^2}{2y^2(\tau)} \tau} \frac{J_{|\nu|}(\mu_n s / y(\tau))}{J_{|\nu|+1}(\mu_n)} ds \right. \\
 & \left. + \frac{1}{2} \int_0^{\tau} y^{\nu+1}(s) \Psi(s) e^{-\frac{\mu_n^2}{2y^2(\tau)} (\tau-s)} \frac{J_{|\nu|}(\mu_n y(s) / y(\tau))}{J_{|\nu|+1}(\mu_n)} ds \right]. \quad (32)
 \end{aligned}$$

Note:

Then the final solution for $u(\tau, z)$ reads

$$u(\tau, z) = 2 \frac{z^{-\nu}}{y^2(\tau)} \sum_{n=1}^{\infty} \left[\int_0^{y(0)} u(0, s) s^{\nu+1} e^{-\frac{\mu_n^2}{2y^2(\tau)} \tau} \frac{J_{|\nu|}(\mu_n s / y(\tau)) J_{|\nu|}(\mu_n z / y(\tau))}{J_{|\nu|+1}^2(\mu_n)} ds \right. \\ \left. + \frac{1}{2} \int_0^{\tau} y^{\nu+1}(s) \Psi(s) e^{-\frac{\mu_n^2}{2y^2(\tau)} (\tau-s)} \frac{J_{|\nu|}(\mu_n y(s) / y(\tau)) J_{|\nu|}(\mu_n z / y(\tau))}{J_{|\nu|+1}^2(\mu_n)} ds \right]. \quad (31)$$

To obtain the equation for Ψ , one needs to differentiate Eq.(31) on z , and then let $z = y(\tau)$. This yields

$$\Psi(\tau) = -\frac{2}{y^{3+\nu}(\tau)} \sum_{n=1}^{\infty} (\mu_n + \nu) \left[\int_0^{y(0)} u(0, s) s^{\nu+1} e^{-\frac{\mu_n^2}{2y^2(\tau)} \tau} \frac{J_{|\nu|}(\mu_n s / y(\tau))}{J_{|\nu|+1}(\mu_n)} ds \right. \\ \left. + \frac{1}{2} \int_0^{\tau} y^{\nu+1}(s) \Psi(s) e^{-\frac{\mu_n^2}{2y^2(\tau)} (\tau-s)} \frac{J_{|\nu|}(\mu_n y(s) / y(\tau))}{J_{|\nu|+1}(\mu_n)} ds \right]. \quad (32)$$

Note: The function $\Psi(\tau)$ can also be found by solving the following Fredholm equation:

$$\int_0^{\infty} e^{-\lambda^2 s / 2} \Psi(s) y^{\nu+1}(s) l_{\nu}(y(s) \lambda) ds = -2 \int_0^{y(0)} q^{\nu+1} l_{\nu}(q \lambda) u(0, q) dq. \quad (33)$$

The problem in half-plane domain $[y(\tau), \infty)$ can be solved via the Weber–Orr transform

$$\bar{u}(\tau, p) = \int_{y(\tau)}^{\infty} z^{\nu+1} W(\tau, p, z) u(\tau, z) dz \quad (34)$$

$$u(\tau, z) = z^{-\nu} \int_0^{\infty} \frac{p W(\tau, p, z)}{V(\tau, p)} \bar{u}(\tau, p) dp.$$

The kernel $W(a, b)$ and the function $V(p)$ are defined as follows, [Bateman and Erdélyi 1953]

$$\begin{aligned} W(\tau, a, b) &= J_{|\nu|}(ab) Y_{|\nu|}(ay(\tau)) - Y_{|\nu|}(ab) J_{|\nu|}(ay(\tau)), \\ V(\tau, p) &= J_{|\nu|}^2(py(\tau)) + Y_{|\nu|}^2(py(\tau)). \end{aligned} \quad (35)$$

The problem in half-plane domain $[y(\tau), \infty)$ can be solved via the Weber–Orr transform

$$\bar{u}(\tau, p) = \int_{y(\tau)}^{\infty} z^{\nu+1} W(\tau, p, z) u(\tau, z) dz \quad (34)$$

$$u(\tau, z) = z^{-\nu} \int_0^{\infty} \frac{p W(\tau, p, z)}{V(\tau, p)} \bar{u}(\tau, p) dp.$$

The kernel $W(a, b)$ and the function $V(p)$ are defined as follows, [Bateman and Erdélyi 1953]

$$\begin{aligned} W(\tau, a, b) &= J_{|\nu|}(ab) Y_{|\nu|}(ay(\tau)) - Y_{|\nu|}(ab) J_{|\nu|}(ay(\tau)), \\ V(\tau, p) &= J_{|\nu|}^2(py(\tau)) + Y_{|\nu|}^2(py(\tau)). \end{aligned} \quad (35)$$

The Webber–Orr transform is a generalization of the Hankel transform to the interval $[a, \infty)$.

The problem in half-plane domain $[y(\tau), \infty)$ can be solved via the Weber–Orr transform

$$\bar{u}(\tau, p) = \int_{y(\tau)}^{\infty} z^{\nu+1} W(\tau, p, z) u(\tau, z) dz \quad (34)$$

$$u(\tau, z) = z^{-\nu} \int_0^{\infty} \frac{p W(\tau, p, z)}{V(\tau, p)} \bar{u}(\tau, p) dp.$$

The kernel $W(a, b)$ and the function $V(p)$ are defined as follows, [Bateman and Erdélyi 1953]

$$\begin{aligned} W(\tau, a, b) &= J_{|\nu|}(ab) Y_{|\nu|}(ay(\tau)) - Y_{|\nu|}(ab) J_{|\nu|}(ay(\tau)), \\ V(\tau, p) &= J_{|\nu|}^2(py(\tau)) + Y_{|\nu|}^2(py(\tau)). \end{aligned} \quad (35)$$

The Webber–Orr transform is a generalization of the Hankel transform to the interval $[a, \infty)$.

The functions $W(\tau, a, b)$ as the functions of the second argument a also form an orthogonal basis in the space $C[y(\tau), \infty)$ for all $\tau > 0$. and the functions $J_{\nu}(xp)$ form an orthogonal basis in the space $C[0, \infty)$.

The problem in half-plane domain $[y(\tau), \infty)$ can be solved via the Weber–Orr transform

$$\bar{u}(\tau, p) = \int_{y(\tau)}^{\infty} z^{\nu+1} W(\tau, p, z) u(\tau, z) dz \quad (34)$$

$$u(\tau, z) = z^{-\nu} \int_0^{\infty} \frac{p W(\tau, p, z)}{V(\tau, p)} \bar{u}(\tau, p) dp.$$

The kernel $W(a, b)$ and the function $V(p)$ are defined as follows, [Bateman and Erdélyi 1953]

$$\begin{aligned} W(\tau, a, b) &= J_{|\nu|}(ab) Y_{|\nu|}(ay(\tau)) - Y_{|\nu|}(ab) J_{|\nu|}(ay(\tau)), \\ V(\tau, p) &= J_{|\nu|}^2(py(\tau)) + Y_{|\nu|}^2(py(\tau)). \end{aligned} \quad (35)$$

The Webber–Orr transform is a generalization of the Hankel transform to the interval $[a, \infty)$.

The functions $W(\tau, a, b)$ as the functions of the second argument a also form an orthogonal basis in the space $C[y(\tau), \infty)$ for all $\tau > 0$. and the functions $J_{\nu}(xp)$ form an orthogonal basis in the space $C[0, \infty)$.

The definitions in Eq.(35) are generalizations of the Pythagorean and Angle sum identities for trigonometric functions to the case of cylinder functions $J_{|\nu|}$ and $Y_{|\nu|}$.

The problem in half-plane domain $[y(\tau), \infty)$ can be solved via the Weber–Orr transform

$$\bar{u}(\tau, p) = \int_{y(\tau)}^{\infty} z^{\nu+1} W(\tau, p, z) u(\tau, z) dz \quad (34)$$

$$u(\tau, z) = z^{-\nu} \int_0^{\infty} \frac{p W(\tau, p, z)}{V(\tau, p)} \bar{u}(\tau, p) dp.$$

The kernel $W(a, b)$ and the function $V(p)$ are defined as follows, [Bateman and Erdélyi 1953]

$$\begin{aligned} W(\tau, a, b) &= J_{|\nu|}(ab) Y_{|\nu|}(ay(\tau)) - Y_{|\nu|}(ab) J_{|\nu|}(ay(\tau)), \\ V(\tau, p) &= J_{|\nu|}^2(py(\tau)) + Y_{|\nu|}^2(py(\tau)). \end{aligned} \quad (35)$$

The Webber–Orr transform is a generalization of the Hankel transform to the interval $[a, \infty)$.

The functions $W(\tau, a, b)$ as the functions of the second argument a also form an orthogonal basis in the space $C[y(\tau), \infty)$ for all $\tau > 0$. and the functions $J_{\nu}(xp)$ form an orthogonal basis in the space $C[0, \infty)$.

The definitions in Eq.(35) are generalizations of the Pythagorean and Angle sum identities for trigonometric functions to the case of cylinder functions $J_{|\nu|}$ and $Y_{|\nu|}$. In particular, for the indexes $\nu = 1/2 + k$, $k \in \mathbb{Z}$ the functions $J_{|\nu|}$ and $Y_{|\nu|}$ can be explicitly represented in terms of sine and cosine functions.

- ▶ We consider Up-and-Out Barrier Call option written on the underlying stock which follows the CEV process with $0 < \beta < 1$. To recall, after the change of variables proposed in this paper is done, the problem transforms to the solution of the Bessel PDE at the domain $\bar{z} \in [y(\tau), \infty)$. Hence, in new variables the Up-and-Out option transforms to the Down-and-Out option.
- ▶ We approach at pricing the Up-and-Out barrier Call option in the CEV model twofold. First, as a benchmark we solve the PDE in Eq.(2) by using a finite-difference (FD) scheme of the second order in space and time. We use the Crank-Nicolson scheme with few first Rannacher steps on a non-uniform grid compressed close to the barrier level, see [Itkin 2017] .

- ▶ We consider Up-and-Out Barrier Call option written on the underlying stock which follows the CEV process with $0 < \beta < 1$. To recall, after the change of variables proposed in this paper is done, the problem transforms to the solution of the Bessel PDE at the domain $\bar{z} \in [y(\tau), \infty)$. Hence, in new variables the Up-and-Out option transforms to the Down-and-Out option.
- ▶ We approach at pricing the Up-and-Out barrier Call option in the CEV model twofold. First, as a benchmark we solve the PDE in Eq.(2) by using a finite-difference (FD) scheme of the second order in space and time. We use the Crank-Nicolson scheme with few first Rannacher steps on a non-uniform grid compressed close to the barrier level, see [Itkin 2017] .
- ▶ Alternatively, we apply the method of Bessel potentials (BP) to solve the Bessel PDE Eq.(5). For doing so, first we solve the Volterra equation in Eq.(19) where the kernel is approximated on a rectangular grid $M \times M$, and the integral is computed using the trapezoidal rule. This implies solving the following system of linear equations

$$\|\zeta\| = (I + P)\|\Psi\|. \quad (36)$$

- ▶ We consider Up-and-Out Barrier Call option written on the underlying stock which follows the CEV process with $0 < \beta < 1$. To recall, after the change of variables proposed in this paper is done, the problem transforms to the solution of the Bessel PDE at the domain $\bar{z} \in [y(\tau), \infty)$. Hence, in new variables the Up-and-Out option transforms to the Down-and-Out option.
- ▶ We approach at pricing the Up-and-Out barrier Call option in the CEV model twofold. First, as a benchmark we solve the PDE in Eq.(2) by using a finite-difference (FD) scheme of the second order in space and time. We use the Crank-Nicolson scheme with few first Rannacher steps on a non-uniform grid compressed close to the barrier level, see [Itkin 2017] .
- ▶ Alternatively, we apply the method of Bessel potentials (BP) to solve the Bessel PDE Eq.(5). For doing so, first we solve the Volterra equation in Eq.(19) where the kernel is approximated on a rectangular grid $M \times M$, and the integral is computed using the trapezoidal rule. This implies solving the following system of linear equations

$$\|\zeta\| = (I + P)\|\Psi\|. \quad (36)$$

- ▶ Here $\|\Psi\|$ is the vector of discrete values of $\Psi(\tau)$, $\tau \in \left[0, \tau(t)\right]_{t=0}$ on a grid with M nodes, $\|\zeta\|$ is a similar vector of $\zeta(\tau)$, I is the unit $M \times M$ matrix, and P is the $M \times M$ matrix of the kernel values on the same grid. Note, that the matrix P is lower triangular. Therefore, solution of Eq.(36) can be done with complexity $O(M^2)$.

- ▶ We consider Up-and-Out Barrier Call option written on the underlying stock which follows the CEV process with $0 < \beta < 1$. To recall, after the change of variables proposed in this paper is done, the problem transforms to the solution of the Bessel PDE at the domain $\bar{z} \in [y(\tau), \infty)$. Hence, in new variables the Up-and-Out option transforms to the Down-and-Out option.
- ▶ We approach at pricing the Up-and-Out barrier Call option in the CEV model twofold. First, as a benchmark we solve the PDE in Eq.(2) by using a finite-difference (FD) scheme of the second order in space and time. We use the Crank-Nicolson scheme with few first Rannacher steps on a non-uniform grid compressed close to the barrier level, see [Itkin 2017] .
- ▶ Alternatively, we apply the method of Bessel potentials (BP) to solve the Bessel PDE Eq.(5). For doing so, first we solve the Volterra equation in Eq.(19) where the kernel is approximated on a rectangular grid $M \times M$, and the integral is computed using the trapezoidal rule. This implies solving the following system of linear equations

$$\|\zeta\| = (I + P)\|\Psi\|. \quad (36)$$

- ▶ Here $\|\Psi\|$ is the vector of discrete values of $\Psi(\tau)$, $\tau \in [0, \tau(t)]_{t=0}$ on a grid with M nodes, $\|\zeta\|$ is a similar vector of $\zeta(\tau)$, I is the unit $M \times M$ matrix, and P is the $M \times M$ matrix of the kernel values on the same grid. Note, that the matrix P is lower triangular. Therefore, solution of Eq.(36) can be done with complexity $O(M^2)$.

In this test we use the explicit form of parameters $r(t), q(t), \sigma(t)$

$$r(t) = r_0 - r_k(a+t), \quad q(t) = q_0 - q_k(a+t), \quad \sigma(t) = \sigma_0\sqrt{a+t}, \quad (37)$$

where $r_0, q_0, \sigma_0, r_k, q_k, \sigma_k$ are constants. We also assume $r_0 = q_0$, and $H = \text{const}$.

In this test we use the explicit form of parameters $r(t), q(t), \sigma(t)$

$$r(t) = r_0 - r_k(a + t), \quad q(t) = q_0 - q_k(a + t), \quad \sigma(t) = \sigma_0 \sqrt{a + t}, \quad (37)$$

where $r_0, q_0, \sigma_0, r_k, q_k, \sigma_k$ are constants. We also assume $r_0 = q_0$, and $H = \text{const}$.

The model parameters for this test are presented in Table 1:

Table 1: Parameters of the test.

r_0	q_0	σ_0	r_k	q_k	σ_k	a	H	S
0.01	0.01	0.3	0.01	0.005	0.2	1.0	100	70

In this test we use the explicit form of parameters $r(t), q(t), \sigma(t)$

$$r(t) = r_0 - r_k(a + t), \quad q(t) = q_0 - q_k(a + t), \quad \sigma(t) = \sigma_0 \sqrt{a + t}, \quad (37)$$

where $r_0, q_0, \sigma_0, r_k, q_k, \sigma_k$ are constants. We also assume $r_0 = q_0$, and $H = \text{const}$.

The model parameters for this test are presented in Table 1:

Table 1: Parameters of the test.

r_0	q_0	σ_0	r_k	q_k	σ_k	a	H	S
0.01	0.01	0.3	0.01	0.005	0.2	1.0	100	70

We run the test for a set of maturities $T \in [1/12, 0.3, 0.5, 1]$ and strikes $K \in [59, 64, 69, 74, 79, 84]$.




Figure 1: Up-and-Out barrier Call option price computed by using the BP method.




Figure 2: % difference of prices computed by using the BP and FD methods.

Table 2: Up-and-Out barrier Call option prices computed by using the BP and FD methods.

	BP				FD				Difference %			
K\T	0.0833	0.3	0.5	1.0	0.0833	0.3	0.5	1.0	0.0833	0.3	0.5	1.0
59	9.3192	3.3642	1.6845	0.4976	9.2924	3.3554	1.6884	0.5175	0.2876	0.2604	-0.2321	-3.9899
64	6.2167	2.1795	1.0671	0.3038	6.2025	2.1831	1.0793	0.3252	0.2286	-0.1654	-1.1438	-7.0291
69	3.8402	1.3219	0.6339	0.1731	3.8341	1.3319	0.6494	0.1931	0.1597	-0.7624	-2.4444	-11.5443
74	2.1608	0.7350	0.3450	0.0891	2.1605	0.7477	0.3606	0.1061	0.0118	-1.7293	-4.5240	-19.1029
79	1.0746	0.3612	0.1652	0.0388	1.0775	0.3736	0.1787	0.0522	-0.2700	-3.4102	-8.1779	-34.6241
84	0.4448	0.1462	0.0641	0.0121	0.4484	0.1561	0.0743	0.0216	-0.7971	-6.7649	-15.9277	-78.1360

- It can be seen that the agreement of both methods is good (less than 1%) if the option price is not too small which occurs when the strike K is close to the barrier or at high maturities. In this case, as this is seen from Table 2, the relative difference becomes large, but the absolute difference of two methods is about one cent, which is almost insignificant. Obviously, such cases are a challenge for any FD method, as at $t = T$ there is a jump in the initial condition at the boundary, and the first derivative of the solution doesn't exist in this point.
- As far as performance of both methods is concerned, to decrease the elapsed time for the FD method instead of Eq.(2) we solve the corresponding forward PDE. Therefore, prices of all options for a given set of strikes and maturities could be obtained within one sweep. This also requires $\bar{m} \times \bar{k}$ integrations of the product of thus found density function with the payoff function, where \bar{m} is the total number of maturities, and \bar{k} is the total number of strikes. In this test the elapsed time for the FD method is, on average, 140 msec.

- ▶ It can be seen that the agreement of both methods is good (less than 1%) if the option price is not too small which occurs when the strike K is close to the barrier or at high maturities. In this case, as this is seen from Table 2, the relative difference becomes large, but the absolute difference of two methods is about one cent, which is almost insignificant. Obviously, such cases are a challenge for any FD method, as at $t = T$ there is a jump in the initial condition at the boundary, and the first derivative of the solution doesn't exist in this point.
- ▶ As far as performance of both methods is concerned, to decrease the elapsed time for the FD method instead of Eq.(2) we solve the corresponding forward PDE. Therefore, prices of all options for a given set of strikes and maturities could be obtained within one sweep. This also requires $\bar{m} \times \bar{k}$ integrations of the product of thus found density function with the payoff function, where \bar{m} is the total number of maturities, and \bar{k} is the total number of strikes. In this test the elapsed time for the FD method is, on average, 140 msec.
- ▶ For the BP method, since the expression for $\zeta(\tau)$ in Eq.(17) is not known in closed form, we compute this integral numerically by using the Simpson quadratures. Nevertheless, to make the results accurate, we need to increase the number of the grid nodes M . As compared with [Itkin and Muravey 2020], where a similar expression for the Hull-White model could be computed in closed form, and $M = 20$ provided a sufficient accuracy, here we need to take $M = 100$. Nevertheless, the elapsed time, on average, is 70 msec, i.e. twice faster than the FD method for the forward equation.

- ▶ It can be seen that the agreement of both methods is good (less than 1%) if the option price is not too small which occurs when the strike K is close to the barrier or at high maturities. In this case, as this is seen from Table 2, the relative difference becomes large, but the absolute difference of two methods is about one cent, which is almost insignificant. Obviously, such cases are a challenge for any FD method, as at $t = T$ there is a jump in the initial condition at the boundary, and the first derivative of the solution doesn't exist in this point.
- ▶ As far as performance of both methods is concerned, to decrease the elapsed time for the FD method instead of Eq.(2) we solve the corresponding forward PDE. Therefore, prices of all options for a given set of strikes and maturities could be obtained within one sweep. This also requires $\bar{m} \times \bar{k}$ integrations of the product of thus found density function with the payoff function, where \bar{m} is the total number of maturities, and \bar{k} is the total number of strikes. In this test the elapsed time for the FD method is, on average, 140 msec.
- ▶ For the BP method, since the expression for $\zeta(\tau)$ in Eq.(17) is not known in closed form, we compute this integral numerically by using the Simpson quadratures. Nevertheless, to make the results accurate, we need to increase the number of the grid nodes M . As compared with [Itkin and Muravey 2020], where a similar expression for the Hull-White model could be computed in closed form, and $M = 20$ provided a sufficient accuracy, here we need to take $M = 100$. Nevertheless, the elapsed time, on average, is 70 msec, i.e. twice faster than the FD method for the forward equation.

Here we present the results of the performance of GIT method. Our numerical scheme is similar to that for the BP method: first we solve the Volterra equation and then compute the value of the integrals using a trapezoidal rule. The comparison with FD benchmark is presented in the following Figure and Table:

Here we present the results of the performance of GIT method. Our numerical scheme is similar to that for the BP method: first we solve the Volterra equation and then compute the value of the integrals using a trapezoidal rule. The comparison with FD benchmark is presented in the following Figure and Table:




Figure 3: Percentage difference of Up-and-Out barrier Call option prices computed by using the FD and GIT methods.

	Difference %			
K\T	0.0833	0.3	0.5	1.0
59	0.71108	-0.2191	-0.3634	-0.7888
64	1.1101	-0.2149	-0.3654	-0.7997
69	1.7729	-0.20803	-0.3556	-0.8132
74	2.6313	-0.1879	-0.3318	-0.8992
79	3.5572	-0.1327	-0.3129	-1.0134
84	4.4718	0.0044	-0.2848	-1.1541

Table 3: Up-and-Out barrier Call option prices computed by using the GIT and FD methods.

- ▶ It can be seen that this method produces very accurate results at high strikes and maturities (i.e. where the option price is relatively small) in contrast to the BP method. The solution obtained by GIT consists of the exponents proportional to the time τ , see [Carr, Itkin, and Muravey 2020] . Contrary, when the price is higher (short maturities, low strikes) the GIT method is slightly less accurate than the BP method, as in Eq.(19) the exponent is inversely proportional to τ .
- ▶ This situation is well investigated for the heat equation with constant coefficients. As applied to pricing double barrier options, it is described in [Lipton 2002] . There exist two representation of the solution: one - obtained by using the method of images, and the other one - by the Fourier series. Despite both solutions are equal as the infinite series, their convergence properties are different. In particular, the Fourier method is superior when the difference between the upper H and lower L barriers is small and the time is relatively large. And the image expansion should be used otherwise.

- ▶ It can be seen that this method produces very accurate results at high strikes and maturities (i.e. where the option price is relatively small) in contrast to the BP method. The solution obtained by GIT consists of the exponents proportional to the time τ , see [Carr, Itkin, and Muravey 2020] . Contrary, when the price is higher (short maturities, low strikes) the GIT method is slightly less accurate than the BP method, as in Eq.(19) the exponent is inversely proportional to τ .
- ▶ This situation is well investigated for the heat equation with constant coefficients. As applied to pricing double barrier options, it is described in [Lipton 2002] . There exist two representation of the solution: one - obtained by using the method of images, and the other one - by the Fourier series. Despite both solutions are equal as the infinite series, their convergence properties are different. In particular, the Fourier method is superior when the difference between the upper H and lower L barriers is small and the time is relatively large. And the image expansion should be used otherwise.
- ▶ In this paper we come to a similar principle for the time-dependent problems, and not just for the heat equation but also for the Bessel one. Thus, it is important that both the BP and GIT methods don't duplicate but rather compliment each other.

- ▶ It can be seen that this method produces very accurate results at high strikes and maturities (i.e. where the option price is relatively small) in contrast to the BP method. The solution obtained by GIT consists of the exponents proportional to the time τ , see [Carr, Itkin, and Muravey 2020] . Contrary, when the price is higher (short maturities, low strikes) the GIT method is slightly less accurate than the BP method, as in Eq.(19) the exponent is inversely proportional to τ .
- ▶ This situation is well investigated for the heat equation with constant coefficients. As applied to pricing double barrier options, it is described in [Lipton 2002] . There exist two representation of the solution: one - obtained by using the method of images, and the other one - by the Fourier series. Despite both solutions are equal as the infinite series, their convergence properties are different. In particular, the Fourier method is superior when the difference between the upper H and lower L barriers is small and the time is relatively large. And the image expansion should be used otherwise.
- ▶ In this paper we come to a similar principle for the time-dependent problems, and not just for the heat equation but also for the Bessel one. Thus, it is important that both the BP and GIT methods don't duplicate but rather compliment each other.
- ▶ It is also worth mentioning that in many situations the parameters of the model are such that the boundary $y(\tau)$ changes slowly with time, i.e. $y(\tau)$ is almost const. Accordingly, one don't need to solve the Volterra equation that makes the algorithm about 2.5 faster.

- ▶ It can be seen that this method produces very accurate results at high strikes and maturities (i.e. where the option price is relatively small) in contrast to the BP method. The solution obtained by GIT consists of the exponents proportional to the time τ , see [Carr, Itkin, and Muravey 2020] . Contrary, when the price is higher (short maturities, low strikes) the GIT method is slightly less accurate than the BP method, as in Eq.(19) the exponent is inversely proportional to τ .
- ▶ This situation is well investigated for the heat equation with constant coefficients. As applied to pricing double barrier options, it is described in [Lipton 2002] . There exist two representation of the solution: one - obtained by using the method of images, and the other one - by the Fourier series. Despite both solutions are equal as the infinite series, their convergence properties are different. In particular, the Fourier method is superior when the difference between the upper H and lower L barriers is small and the time is relatively large. And the image expansion should be used otherwise.
- ▶ In this paper we come to a similar principle for the time-dependent problems, and not just for the heat equation but also for the Bessel one. Thus, it is important that both the BP and GIT methods don't duplicate but rather compliment each other.
- ▶ It is also worth mentioning that in many situations the parameters of the model are such that the boundary $y(\tau)$ changes slowly with time, i.e. $y(\tau)$ is almost const. Accordingly, one don't need to solve the Volterra equation that makes the algorithm about 2.5 faster.



Abramowitz, M. and I. Stegun (1964). *Handbook of Mathematical Functions*. Dover Publications, Inc.



Andersen, L.B.G. and V.V. Piterbarg (2010). *Interest Rate Modeling*. Interest Rate Modeling v. 2. Atlantic Financial Press. ISBN: 9780984422111.



Bateman, H. and A. Erdélyi (1953). *Higher Transcendental Functions*. Vol. 1. Bateman Manuscript Project California Institute of Technology. McGraw-Hill.



Carr, P., A. Itkin, and D. Muravey (2020). "Semi-closed form prices of barrier options in the time-dependent CEV and CIR models". In: *Journal of Derivatives* 28.1, pp. 26–50.



Carr, P. and V. Linetsky (2006). "A jump to default extended CEV model: an application of Bessel processes". In: *Finance and Stochastics* 10, pp. 303–330.



Cox, J. (1975). *Notes on Option Pricing I. Constant Elasticity of Variance Diffusions*. Tech. rep. Stanford University working paper.



Emanuel, D. and J. Macbeth (1982). "Further Results on the Constant Elasticity of Variance Call Option Pricing Model". In: *Journal of Financial and Quantitative Analysis* 17, pp. 533–554.



Itkin, A. (2017). *Pricing Derivatives Under Lévy Models. Modern Finite-Difference and Pseudo-Differential Operators Approach*. Vol. 12. Pseudo-Differential Operators. Birkhauser.



Itkin, A. and D. Muravey (Apr. 2020). *Semi-closed form prices of barrier options in the Hull-White model*. Arxiv:2004.09591.



Lawler, Gregory F. (2018). "Notes on the Bessel Process". In: Corpus ID: 52200396. URL: <http://www.math.uchicago.edu/~lawler/bessel18new.pdf>.



Linetsky, V. and R. Mendoza (2010). "Encyclopedia of Quantitative Finance". In: *Constant Elasticity of Variance (CEV) Diffusion Model*. John Wiley & Sons. ISBN: 9780470061602.



Lipton, A. (2002). "The vol smile problem". In: *Risk*, pp. 61–65.



Polyanin, A.D. (2002). *Handbook of linear partial differential equations for engineers and scientists*. Chapman & Hall/CRC.



Polyanin, P. and A.V. Manzhirov (2008). *Handbook of Integral Equations: Second Edition*. Handbooks of mathematical equations. CRC Press. ISBN: 9780203881057.



Revuz, D. and M. Yor (1999). *Continuous Martingales and Brownian Motion*. 3rd. Berlin, Germany: Springer.



Tikhonov, A.N. and A.A. Samarskii (1963). *Equations of mathematical physics*. Oxford: Pergamon Press.

Thank you!